

For research use only




## Anti Human Macrophage Scavenger Receptor A (MSR- A:CD204) Monoclonal Antibody (Clone No. SRA-C6)

Class A macrophage scavenger receptor (MSR-A: CD204) was identified in the search for the receptor molecules that are implicated in the pathological deposition of cholesterol during atherogenesis through receptor-mediated uptake of modified low density lipoprotein (LDL). MSR-A possesses a wide range of ligand-binding specificities and recognize a variety of molecules such as modified LDL including acetylated LDL, oxidized LDL, advanced glycation end products (AGEs), polyribonucleotides such as poly G and poly I and bacterial surface lipids including lipopolysaccharide and lipoteichoic acid.

This antibody was produced from the mouse immunized with recombinant protein of human type I MSR-A, and has been proved to be useful for the western blotting and immunohistochemistry. This antibody also inhibits the endocytic degradation of acetylated LDL and oxidized LDL by high glucose-treated human monocyte-derived macrophages and has anti MSR-A neutralizing activity.

This antibody is useful tools for the study of MSR-A in atherogenesis and various other pathological conditions.

Package Size	50 $\mu$ g (200 $\mu$ l / vial)
Format	Mouse monoclonal antibody 0.25mg/ml
Buffer	PBS [containing 2% Block Ace as a stabilizer, 0.1%Proclin as a bacteriostat]
Storage	Store below -20°C
	Once thawed, store at 4°C. Repeated freeze-thaw cycles should be avoided.
Clone No.	SRA-C6
Subclass	IgG1
Purification method	The spleen cells obtained from MSR-A deficient mouse, immunized with recombinant protein corresponding to amino acid 131-451 of human type I MSR-A, were fused with mouse NS-1 myeloma cells. The hybridoma cell line with positive reaction was grown on non-serum medium, from which the antibody was purified by Protein G affinity chromatography.
Working dilution for	immunohistochemistry: 5.0 $\mu$ g/ml, western blotting : 1.0 $\mu$ g/ml, Neutralization: Depends on the experimental design(Application Reference:1)



**Left:** Human alveolar macrophages(Cytospin preparation): Most macrophages are positive.

**Center:** Human liver (paraffin section): Kupffer cells are positive

**Right:** Neutralizing activity of SRA-C6 (20  $\mu$ g/ml): Inhibitory effect of anti-human SR-A antibody on the degradation of <sup>125</sup>I-AcLDL by human monocyte-derived macrophages(day7)

Takeya M., Department of Cell Pathology, Graduate School of Medical Sciences, Kumamoto University, Kumamoto, Japan

## Anti Human Macrophage Scavenger Receptor A (MSR- A:CD204) Monoclonal Antibody (Clone No. SRA-C6)

### 【Specificity】

Organ	Reaction		Organ	Reaction	
	Positive	Negative		Positive	Negative
Heart	Intramuscular M $\phi$ (+-)		Trachea	Mucosal M $\phi$ (+-)	
Lung	Alveolar M $\phi$ (+) M $\phi$ in alveolar septa (+-)		Esophagus	Interstitial M $\phi$ (+-)	
Liver	Kupffer cells (+) M $\phi$ in portal triads(+)		Stomach	M $\phi$ in lamina propria(+) M $\phi$ in striated muscle(+/-)	
Kidney	Interstitial M $\phi$ (+)		Small and large intestines	M $\phi$ in lamina propria(+) M $\phi$ in striated muscle(+/-)	
Spleen	Red pulp M $\phi$ (+)	Interdigitating cells	Skin	Dermal M $\phi$ (+)	Langerhans cells
Thymus	Interlobular M $\phi$ (+)		Brain (cerebrum and cerebellum)	Perivascular M $\phi$ (Mato cells) (+)	
Lymph nodes	Sinus M $\phi$ (+)	Tingible body M $\phi$ Interdigitating cells	Testes	Interstitial M $\phi$ (+)	
Pancreas	Interlobular M $\phi$ (+)		Uterus	Interstitial M $\phi$ (+)	
Salivary gland	Interlobular M $\phi$ (+)		Ovaries	Interstitial M $\phi$ (+)	
Thyroid	Interfollicular M $\phi$ (+-)		Placenta	Hofbauer cells (+)	
Parathyroid	Interlobular M $\phi$ (+-)		Bone marrow	M $\phi$ (+)	Myeloid precursor cells
Adrenals	Interstitial M $\phi$ (+)		Blood monocyte	3 days in culture (+)	Freshly isolated
Urinary bladder	Interstitial M $\phi$ (+-)				
Prostate	Interstitial M $\phi$ (+-)				

M  $\phi$  : macrophage 、 (+): most cells were positive; (+-): about 10-50% of cells were positive

### 【Application Reference】

1. Fukuhara-Takaki K., Sakai M., Sakamoto Y., Takeya M., Horiuchi S.: Expression of class A scavenger receptor is enhanced by high glucose in vitro and under diabetic conditions in vivo: one mechanism for an increased rate of atherosclerosis in diabetes.: J Biol Chem. 280(5): 3355-3364, 2005
2. Tomokiyo R., Jinnouchi K., Honda M., Wada Y., Hanada N., Hiraoka T., Suzuki H., Kodama T., Takahashi K., Takeya M.: Production, characterization, and interspecies reactivities of monoclonal antibodies against human class A macrophage scavenger receptors: Atherosclerosis, 161:123-132, 2002

#### Manufacturer



7-1-14 Minatojimaminami-machi, Chuo-ku, Kobe, Japan 650-0047

Telephone: +81-78-306-0295 FAX:+81-78-306-0296

URL:<http://www.transgenic.co.jp> techstaff@transgenic.co.jp